

Dibromido[2-[(4-fluorophenyl)imino-methyl]pyridine- κ^2N,N']zinc

Saeed Dehghanpour* and Ali Mahmoudi

Department of Chemistry, Islamic Azad University, Karaj, Iran
Correspondence e-mail: Dehghanpour_farasha@yahoo.com


Received 30 June 2012; accepted 8 July 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.030; wR factor = 0.072; data-to-parameter ratio = 18.1.

In the title complex, $[\text{ZnBr}_2(\text{C}_{12}\text{H}_9\text{FN}_2)]$, the Zn^{II} atom has a distorted tetrahedral Br_2N_2 coordination sphere. The organic ligand is bidentate, coordinating the Zn^{II} atom via two imine N atoms. The benzene and pyridine rings are oriented at a dihedral angle of 10.49 (1)°. In the crystal, weak $\text{C}-\text{H}\cdots\text{F}$ and $\text{C}-\text{H}\cdots\text{Br}$ hydrogen bonds are observed.

Related literature

For background information, see: Dehghanpour *et al.* (2009). For related structures, see: Dehghanpour *et al.* (2007); Salehzadeh *et al.* (2011); Khalaj *et al.* (2009).



Experimental

Crystal data

$[\text{ZnBr}_2(\text{C}_{12}\text{H}_9\text{FN}_2)]$
 $M_r = 425.40$
Monoclinic, $P2_1/c$
 $a = 7.7351$ (10) Å
 $b = 9.5372$ (13) Å
 $c = 18.501$ (2) Å
 $\beta = 96.052$ (3)°
 $V = 1357.2$ (3) Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 7.69$ mm⁻¹
 $T = 100$ K

$0.17 \times 0.06 \times 0.04$ mm

Data collection

Bruker APEXII area-detector diffractometer
Absorption correction: multi-scan (APEX2; Bruker, 2005)
 $T_{\min} = 0.435$, $T_{\max} = 0.734$
18332 measured reflections
2956 independent reflections
2469 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.072$
 $S = 1.00$
2956 reflections
163 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 2.44$ e Å⁻³
 $\Delta\rho_{\min} = -0.77$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-H\cdots A$ | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|---|-------|-------------|-------------|---------------|
| $\text{C12}-\text{H12A}\cdots\text{Br1}$ | 0.93 | 3.04 | 3.866 (4) | 148 |
| $\text{C2}-\text{H2A}\cdots\text{F1}^{\text{i}}$ | 0.93 | 2.50 | 3.081 (4) | 121 |
| $\text{C5}-\text{H5A}\cdots\text{Br1}^{\text{ii}}$ | 0.93 | 3.01 | 3.767 (4) | 140 |
| $\text{C6}-\text{H6A}\cdots\text{Br1}^{\text{iii}}$ | 0.93 | 3.05 | 3.756 (4) | 134 |
| $\text{C3}-\text{H3A}\cdots\text{Br2}^{\text{iv}}$ | 0.93 | 2.90 | 3.810 (4) | 166 |

Symmetry codes: (i) $x, -y + \frac{5}{2}, z - \frac{1}{2}$; (ii) $-x, -y + 1, -z$; (iii) $-x, -y + 2, -z$; (iv) $x, -y + \frac{3}{2}, z - \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors would like to acknowledge the Islamic Azad University Research Councils for partial support of this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2564).

References

- Bruker (2005). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dehghanpour, S., Khalaj, M. & Mahmoudi, A. (2009). *Polyhedron*, **28**, 1205–1210.
- Dehghanpour, S., Mahmoudi, A., Khalaj, M. & Salmanpour, S. (2007). *Acta Cryst. E* **63**, m2840.
- Khalaj, M., Dehghanpour, S., Mahmoudi, A. & Seyedidarzam, S. (2009). *Acta Cryst. E* **65**, m890.
- Salehzadeh, S., Khalaj, M., Dehghanpour, S. & Tarmoradi, I. (2011). *Acta Cryst. E* **67**, m1556.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, m1113 [doi:10.1107/S1600536812031091]

Dibromido{2-[(4-fluorophenyl)iminomethyl]pyridine- κ^2N,N' }zinc

Saeed Dehghanpour and Ali Mahmoudi

Comment

In continuation of our research on the synthesis and characterization of metal complexes containing bidentate Schiff base ligands (Dehghanpour *et al.* (2009)), we now report the synthesis and crystal structure of a zinc complex of the Schiff base, (4-fluorophenyl)pyridin-2-ylmethyleneamine.

The metal centre in the title complex (Fig. 1) has a tetrahedral coordination which shows significant distortion, mainly due to the presence of the five-membered chelate ring. The endocyclic N1—Zn1—N2 angle (81.13 (12)°) is much narrower than the ideal tetrahedral angle of 109.5°, whereas the opposite Br1—Zn1—Br2 angle (116.72 (2)°) is much wider than the ideal tetrahedral angle. The Zn—Br and Zn—N bond distances compare well with the values found in other tetrahedral Schiff base adducts of zinc bromide (Salehzadeh *et al.*, 2011; Dehghanpour *et al.*, 2007; Khalaj *et al.*, 2009). The interplanar angles between the benzene and pyridine rings in the title structure is 10.49 (1)°. In the crystal, weak C—H \cdots F and C—H \cdots Br hydrogen bonds are also observed (Tab. 1 & Fig. 2).

Experimental

The title complex was prepared by the reaction of ZnBr₂ (22.5 mg, 0.1 mmol) and (4-fluorophenyl)pyridin-2-ylmethyleneamine (20 mg, 0.1 mmol) in 10 ml of methanol at room temperature. The solution was allowed to stand at room temperature and crystals of the title compound suitable for X-ray analysis formed within a few days.

Refinement

Though the H-atoms were observable in the difference electron density maps they were included at geometrically idealized positions with C—H distances = 0.93 Å and $U_{iso} = 1.2$ times U_{eq} of the atoms to which they were bonded. There is a high positive residual density of 2.44 e Å⁻³ near the Zn1 center due to considerable absorption effects which could not be completely corrected.

Computing details

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2* (Bruker, 2005); data reduction: *APEX2* (Bruker, 2005); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).




Figure 1

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.




Figure 2

A view of the C—H⋯F and C—H⋯Br hydrogen bonds (dotted lines) in the crystal structure of the title compound. H atoms non-participating in hydrogen-bonding were omitted for clarity.

Dibromido{2-[(4-fluorophenyl)iminomethyl]pyridine- κ^2N,N' }zinc

Crystal data

[ZnBr₂(C₁₂H₉FN₂)]

M_r = 425.40

Monoclinic, *P*2₁/*c*

Hall symbol: -*P* 2₁/c

a = 7.7351 (10) Å

b = 9.5372 (13) Å

c = 18.501 (2) Å

β = 96.052 (3)°

V = 1357.2 (3) Å³

Z = 4

$F(000) = 816$
 $D_x = 2.082 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 3858 reflections
 $\theta = 2.2\text{--}29.7^\circ$

$\mu = 7.69 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
Prism, colourless
 $0.17 \times 0.06 \times 0.04 \text{ mm}$

Data collection

Bruker APEXII area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(APEX2; Bruker, 2005)
 $T_{\min} = 0.435$, $T_{\max} = 0.734$

18332 measured reflections
2956 independent reflections
2469 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.049$
 $\theta_{\max} = 27.0^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -9 \rightarrow 9$
 $k = -12 \rightarrow 12$
 $l = -23 \rightarrow 23$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.072$
 $S = 1.00$
2956 reflections
163 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0358P)^2 + 2.3621P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 2.44 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.77 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|-------------|---------------|----------------------------------|
| Br1 | −0.00870 (4) | 0.67594 (4) | 0.09661 (2) | 0.01923 (10) |
| Br2 | 0.50834 (4) | 0.69666 (4) | 0.11656 (2) | 0.02040 (10) |
| Zn1 | 0.24082 (5) | 0.76510 (4) | 0.05390 (2) | 0.01606 (11) |
| N2 | 0.2404 (4) | 0.9803 (3) | 0.03443 (16) | 0.0159 (6) |
| C1 | 0.2874 (4) | 0.8917 (4) | −0.08143 (19) | 0.0166 (7) |
| F1 | 0.1273 (3) | 1.3856 (3) | 0.23727 (12) | 0.0320 (6) |
| C8 | 0.2592 (5) | 1.2286 (4) | 0.0745 (2) | 0.0193 (8) |
| H8A | 0.3069 | 1.2546 | 0.0324 | 0.023* |
| C9 | 0.2290 (5) | 1.3290 (4) | 0.1259 (2) | 0.0227 (8) |
| H9A | 0.2547 | 1.4229 | 0.1187 | 0.027* |
| C2 | 0.3253 (5) | 0.9156 (4) | −0.1518 (2) | 0.0223 (8) |
| H2A | 0.3411 | 1.0063 | −0.1683 | 0.027* |

| | | | | |
|------|------------|------------|---------------|------------|
| C10 | 0.1598 (5) | 1.2862 (4) | 0.1881 (2) | 0.0232 (8) |
| C6 | 0.2678 (4) | 1.0072 (4) | −0.03110 (19) | 0.0180 (7) |
| H6A | 0.2753 | 1.0996 | −0.0466 | 0.022* |
| C12 | 0.1525 (5) | 1.0492 (4) | 0.1500 (2) | 0.0205 (8) |
| H12A | 0.1290 | 0.9552 | 0.1582 | 0.025* |
| C4 | 0.3159 (5) | 0.6678 (4) | −0.1709 (2) | 0.0259 (9) |
| H4A | 0.3237 | 0.5900 | −0.2006 | 0.031* |
| C7 | 0.2183 (4) | 1.0901 (4) | 0.08600 (19) | 0.0160 (7) |
| C3 | 0.3393 (5) | 0.8013 (5) | −0.1973 (2) | 0.0264 (9) |
| H3A | 0.3642 | 0.8144 | −0.2450 | 0.032* |
| C5 | 0.2804 (5) | 0.6514 (4) | −0.0995 (2) | 0.0232 (8) |
| H5A | 0.2662 | 0.5615 | −0.0817 | 0.028* |
| C11 | 0.1222 (5) | 1.1492 (4) | 0.2015 (2) | 0.0240 (8) |
| H11A | 0.0771 | 1.1237 | 0.2443 | 0.029* |
| N1 | 0.2661 (4) | 0.7613 (3) | −0.05549 (16) | 0.0176 (6) |

Atomic displacement parameters (Å²)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|--------------|--------------|-------------|---------------|--------------|--------------|
| Br1 | 0.01582 (18) | 0.01869 (19) | 0.0241 (2) | −0.00193 (13) | 0.00645 (13) | 0.00462 (15) |
| Br2 | 0.01554 (18) | 0.0241 (2) | 0.0222 (2) | 0.00173 (14) | 0.00489 (13) | 0.00680 (15) |
| Zn1 | 0.0172 (2) | 0.0141 (2) | 0.0177 (2) | −0.00006 (16) | 0.00594 (15) | 0.00254 (16) |
| N2 | 0.0161 (15) | 0.0135 (15) | 0.0183 (15) | −0.0001 (12) | 0.0029 (11) | −0.0010 (12) |
| C1 | 0.0130 (16) | 0.0193 (19) | 0.0175 (17) | 0.0004 (14) | 0.0011 (13) | 0.0007 (15) |
| F1 | 0.0414 (14) | 0.0251 (13) | 0.0298 (13) | 0.0036 (11) | 0.0046 (10) | −0.0122 (10) |
| C8 | 0.0170 (17) | 0.0229 (19) | 0.0182 (18) | 0.0014 (15) | 0.0020 (14) | 0.0022 (15) |
| C9 | 0.026 (2) | 0.0112 (18) | 0.030 (2) | 0.0010 (14) | −0.0023 (16) | −0.0023 (16) |
| C2 | 0.0223 (19) | 0.022 (2) | 0.023 (2) | 0.0031 (15) | 0.0046 (15) | 0.0059 (16) |
| C10 | 0.0210 (19) | 0.025 (2) | 0.0231 (19) | 0.0029 (16) | −0.0008 (15) | −0.0089 (17) |
| C6 | 0.0142 (17) | 0.0202 (19) | 0.0198 (18) | 0.0021 (14) | 0.0026 (13) | 0.0039 (15) |
| C12 | 0.0218 (19) | 0.0165 (18) | 0.0238 (19) | −0.0012 (15) | 0.0058 (15) | 0.0024 (15) |
| C4 | 0.028 (2) | 0.027 (2) | 0.023 (2) | 0.0038 (17) | 0.0005 (16) | −0.0125 (17) |
| C7 | 0.0113 (16) | 0.0175 (18) | 0.0192 (18) | 0.0041 (13) | 0.0016 (13) | −0.0037 (14) |
| C3 | 0.028 (2) | 0.036 (2) | 0.0157 (18) | 0.0081 (18) | 0.0056 (15) | 0.0008 (17) |
| C5 | 0.0207 (19) | 0.020 (2) | 0.030 (2) | −0.0019 (15) | 0.0052 (16) | −0.0001 (16) |
| C11 | 0.026 (2) | 0.028 (2) | 0.0185 (19) | 0.0001 (16) | 0.0077 (15) | −0.0019 (16) |
| N1 | 0.0155 (15) | 0.0181 (16) | 0.0198 (15) | 0.0003 (12) | 0.0045 (12) | −0.0007 (13) |

Geometric parameters (Å, °)

| | | | |
|---------|------------|----------|-----------|
| Br1—Zn1 | 2.3225 (6) | C2—C3 | 1.389 (6) |
| Br2—Zn1 | 2.3550 (6) | C2—H2A | 0.9300 |
| Zn1—N1 | 2.054 (3) | C10—C11 | 1.366 (6) |
| Zn1—N2 | 2.084 (3) | C6—H6A | 0.9300 |
| N2—C6 | 1.279 (5) | C12—C11 | 1.385 (5) |
| N2—C7 | 1.439 (4) | C12—C7 | 1.393 (5) |
| C1—N1 | 1.350 (5) | C12—H12A | 0.9300 |
| C1—C2 | 1.383 (5) | C4—C3 | 1.383 (6) |
| C1—C6 | 1.461 (5) | C4—C5 | 1.385 (6) |
| F1—C10 | 1.356 (4) | C4—H4A | 0.9300 |

| | | | |
|---------------|------------|----------------|------------|
| C8—C7 | 1.380 (5) | C3—H3A | 0.9300 |
| C8—C9 | 1.386 (5) | C5—N1 | 1.339 (5) |
| C8—H8A | 0.9300 | C5—H5A | 0.9300 |
| C9—C10 | 1.381 (6) | C11—H11A | 0.9300 |
| C9—H9A | 0.9300 | | |
| N1—Zn1—N2 | 81.13 (12) | N2—C6—C1 | 119.4 (3) |
| N1—Zn1—Br1 | 119.84 (8) | N2—C6—H6A | 120.3 |
| N2—Zn1—Br1 | 115.66 (8) | C1—C6—H6A | 120.3 |
| N1—Zn1—Br2 | 108.07 (8) | C11—C12—C7 | 119.7 (4) |
| N2—Zn1—Br2 | 110.09 (8) | C11—C12—H12A | 120.1 |
| Br1—Zn1—Br2 | 116.72 (2) | C7—C12—H12A | 120.1 |
| C6—N2—C7 | 121.7 (3) | C3—C4—C5 | 119.2 (4) |
| C6—N2—Zn1 | 111.4 (3) | C3—C4—H4A | 120.4 |
| C7—N2—Zn1 | 126.9 (2) | C5—C4—H4A | 120.4 |
| N1—C1—C2 | 122.1 (3) | C8—C7—C12 | 120.6 (3) |
| N1—C1—C6 | 116.3 (3) | C8—C7—N2 | 123.3 (3) |
| C2—C1—C6 | 121.5 (3) | C12—C7—N2 | 116.1 (3) |
| C7—C8—C9 | 119.8 (3) | C4—C3—C2 | 119.1 (4) |
| C7—C8—H8A | 120.1 | C4—C3—H3A | 120.5 |
| C9—C8—H8A | 120.1 | C2—C3—H3A | 120.5 |
| C10—C9—C8 | 118.4 (4) | N1—C5—C4 | 121.9 (4) |
| C10—C9—H9A | 120.8 | N1—C5—H5A | 119.0 |
| C8—C9—H9A | 120.8 | C4—C5—H5A | 119.0 |
| C1—C2—C3 | 118.7 (4) | C10—C11—C12 | 118.6 (4) |
| C1—C2—H2A | 120.7 | C10—C11—H11A | 120.7 |
| C3—C2—H2A | 120.7 | C12—C11—H11A | 120.7 |
| F1—C10—C11 | 119.3 (4) | C5—N1—C1 | 118.9 (3) |
| F1—C10—C9 | 117.9 (3) | C5—N1—Zn1 | 129.5 (3) |
| C11—C10—C9 | 122.8 (4) | C1—N1—Zn1 | 111.2 (2) |
| N1—Zn1—N2—C6 | −4.7 (3) | C6—N2—C7—C12 | 166.6 (3) |
| Br1—Zn1—N2—C6 | −123.7 (2) | Zn1—N2—C7—C12 | −15.4 (4) |
| Br2—Zn1—N2—C6 | 101.4 (2) | C5—C4—C3—C2 | −0.6 (6) |
| N1—Zn1—N2—C7 | 177.1 (3) | C1—C2—C3—C4 | −0.3 (6) |
| Br1—Zn1—N2—C7 | 58.2 (3) | C3—C4—C5—N1 | 0.8 (6) |
| Br2—Zn1—N2—C7 | −76.8 (3) | F1—C10—C11—C12 | 178.6 (3) |
| C7—C8—C9—C10 | 0.7 (5) | C9—C10—C11—C12 | −0.9 (6) |
| N1—C1—C2—C3 | 0.9 (5) | C7—C12—C11—C10 | −0.7 (6) |
| C6—C1—C2—C3 | −178.9 (3) | C4—C5—N1—C1 | −0.2 (5) |
| C8—C9—C10—F1 | −178.6 (3) | C4—C5—N1—Zn1 | −172.7 (3) |
| C8—C9—C10—C11 | 0.9 (6) | C2—C1—N1—C5 | −0.7 (5) |
| C7—N2—C6—C1 | −179.3 (3) | C6—C1—N1—C5 | 179.2 (3) |
| Zn1—N2—C6—C1 | 2.4 (4) | C2—C1—N1—Zn1 | 173.1 (3) |
| N1—C1—C6—N2 | 3.2 (5) | C6—C1—N1—Zn1 | −7.0 (4) |
| C2—C1—C6—N2 | −176.9 (3) | N2—Zn1—N1—C5 | 179.3 (3) |
| C9—C8—C7—C12 | −2.3 (5) | Br1—Zn1—N1—C5 | −66.1 (3) |
| C9—C8—C7—N2 | 178.0 (3) | Br2—Zn1—N1—C5 | 71.0 (3) |
| C11—C12—C7—C8 | 2.3 (5) | N2—Zn1—N1—C1 | 6.3 (2) |

| | | | |
|---------------|------------|---------------|------------|
| C11—C12—C7—N2 | −178.0 (3) | Br1—Zn1—N1—C1 | 120.9 (2) |
| C6—N2—C7—C8 | −13.7 (5) | Br2—Zn1—N1—C1 | −102.0 (2) |
| Zn1—N2—C7—C8 | 164.3 (3) | | |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|--|-------------|---------------------|----------------------------|-------------------------------|
| C12—H12 <i>A</i> \cdots Br1 | 0.93 | 3.04 | 3.866 (4) | 148 |
| C2—H2 <i>A</i> \cdots F1 ⁱ | 0.93 | 2.50 | 3.081 (4) | 121 |
| C5—H5 <i>A</i> \cdots Br1 ⁱⁱ | 0.93 | 3.01 | 3.767 (4) | 140 |
| C6—H6 <i>A</i> \cdots Br1 ⁱⁱⁱ | 0.93 | 3.05 | 3.756 (4) | 134 |
| C3—H3 <i>A</i> \cdots Br2 ^{iv} | 0.93 | 2.90 | 3.810 (4) | 166 |

Symmetry codes: (i) $x, -y+5/2, z-1/2$; (ii) $-x, -y+1, -z$; (iii) $-x, -y+2, -z$; (iv) $x, -y+3/2, z-1/2$.